

CHERRY PESTICIDES – 2010



This sheet lists some pesticides used on cherries. The health symptoms listed may be caused by things other than pesticides. If you feel sick after exposure to pesticides, call a poison control center or doctor for treatment advice. If a person is not breathing, call 911 or an ambulance, give artificial respiration.



Pesticide (ex. Brand Name) Highly Toxic 	Product Use Changes Read the Label (Reentry intervals may vary by product) DO NOT ENTER TREATED AREAS FOR:	Early Symptoms of Acute Pesticide Poisoning
 Azinphos methyl (Guthion)	15 days	  
Carbaryl (Sevin)	12 hours	  
Chlorpyrifos	4 days	  
Dimethoate	10 days/14 days	
Malathion	12 hours	
 Copper Hydroxide	24 hours	   
Myclobutanil (Rally)	24 hours	 
 Paraquat (Firestorm, Gramoxone)	24 hours	   
If you are exposed to a pesticide spray or drift:		If you feel sick while working with pesticides:
<ul style="list-style-type: none"> • Leave the area • Tell your employer • Immediately rinse skin with plenty of water for 15-20 minutes • If you get pesticides in your eyes, hold eyes open and rinse slowly and gently with water for 15-20 minutes. If you wear contacts, rinse for 5 minutes, remove contacts and continue rinsing eyes. • Change into clean clothes • Always wash work clothes separately from other clothes 		<ul style="list-style-type: none"> • Leave the area • Tell your employer you feel sick, start first aid if needed • Go to a doctor or clinic • Have someone go with you • Tell the doctor the pesticide name. Your employer must provide this information <p>Phone numbers:</p> <p>WSDA toll free: 1-877-301-4555 L&I toll free: 1-800-423-7233 DOH (Olympia): 360-236-3360 Poison Emergency 1-800-222-1222</p>