

April 1, 2003

## 2002 Pheromone-trap Detection Survey for Leek Moth, *Acrolepiopsis assectella* (Zeller, 1893) (Lepidoptera: Acrolepiidae), an Exotic Pest of *Allium* spp.

Eric H. LaGasa<sup>1</sup>, John Agnesani<sup>2</sup>, Sahabra Tipton<sup>2</sup>, and Denise Bowden<sup>2</sup>

### Background

Native to Asia, the Leek Moth (LM) was first found in North America in 1993, in Ottawa, Canada, where the pest is now established (CFIA, 2001). To date, the pest has not been detected in the United States (Figure 1 - APHIS NAPIS, 2003).

In Europe, the pest causes heavy damage to leeks, onions, garlic, and related crops by mining and feeding within the foliage and bulbs (see Figure 2), and infested bulbs are subject to extensive rotting in storage. Larval damage to stalks can also reduce seed production (USDA-ARS, 1960). Pest status is characterized as a “minor pest of onions” in some references (Hill, 1983 and 1987), a “serious pest in continental Europe” (Carter, 1984), and a potential pest of native (North American) and ornamental plants in the genus *Allium* (APHIS NPAG, 2000).

### 2002 Project Objective

#### Conduct pheromone-trap detection survey throughout populous western Washington.

- Place and monitor pheromone-traps in areas of commercial and home garden *Allium* culture.
- Screen and identify captured specimens, including non-target material when possible.

### Project Methods and Materials

Three hundred and ninety-six pheromone-lure baited traps were placed in counties along the Interstate-5 corridor in western Washington, from the Canadian border south to Clark County on the Columbia River / Oregon border. Trap placement, by county, is presented in Table 1. Traps were hung in roadside or residential yard trees, primarily in areas where home gardens could provide *Allium* spp. hosts (ornamental and table crop).

Trap placement began in June to allow for completion of all initial trap sets by the beginning of expected adult moth flight in late-June or early-July, and most traps were removed by the end of August.

Figure 1. Leek Moth Survey in U.S., 2003 (USDA)




Figure 2. Leek Moth Life Stages and Damage



<sup>1</sup>Chief Entomologist - Washington State Dept. of Agriculture, Olympia, Washington 98504-2560

<sup>2</sup>Entomology Aides (Project) - Washington State Dept. of Agriculture, Olympia, Washington 98504-2560

**Project Methods and Materials (Cont.)**

Pherocon 2® type traps (a.k.a. "diamond" traps) were used in this survey, based on trap performance in prior WSDA CAPS surveys, ease of use, and the avoidance of small-bird capture/mortality (which is a problem with other trap designs, particularly "wing-traps"). Traps were baited with pheromone-lures provided by the USDA APHIS Otis Methods Development Center. The leek moth pheromone-lures consisted of gray rubber septa (West Co., Lionville, PA. cat. no. 1060-0275), each loaded with 0.1 ml (1 mg) of (Z)-11-Hexadecanol.

Pheromone lures were changed every two weeks as much as possible during the expected (probable) period of adult flight. Traps with specimens were processed at the Olympia Entomology Lab, where captured specimens were identified and counted. Selected specimens were removed from the traps with Hemo-D citrus based solvent and had genitalia extracted and cleared (in KOH) for identification.

**Project Results**

**No leek moth specimens were collected in this survey.**

**Pertinent Literature**

Alford, D. 1999. *A Textbook of Agricultural Entomology*. Blackwell Science.

APHIS NPAG, 2000. USDA New Pest Advisory Group Data: *A. assectella*, Leek Moth – Potential Threat to American Onion and *Allium* Growers. August 1, 2000

Carter, D. 1984. *Pest Lepidoptera of Europe*. Dr. W. Junk, Boston  
CFIA, 2001 (Web Document) Canadian Food Inspection Agency - Plant Pest Info. Acrolepiopsis assectella (Zeller, 1839) Leek Moth <http://www.inspection.gc.ca/english/ppc/science/pps/datasheets/acrasse.shtml> last modified February 14, 2003

Hill, D. 1983. *Agricultural Insect Pests of the Tropics and Their Control*. Second edition. Cambridge University Press, New York.

Hill, D. 1987. *Agricultural Insect Pests of Temperate Regions and Their Control*. Cambridge University Press, New York.

USDA-ARS, 1960. Leek Moth (*A. assectella* Zell.) *Insects Not Known to Occur in the United States* 10:33-34

Table 1. Leek Moth Trap Placement

County	Number of Leek Moth Trap Sites
Whatcom	98
Snohomish	11
Skagit	40
King	75
Pierce	51
Thurston	52
Lewis	12
Cowlitz	21
Clark	36
<b>Total Traps</b>	<b>396</b>

Table 2. Non-target Pheromone-trap Captures

Defoliator species	Number of specimens
<i>Acleris variegana</i>	3
<i>Agonopterix alstroemeriana</i>	1
<i>Apotomis spurinifida</i>	27
<i>Archips fuscocupreanus</i>	4
<i>Archips rosanus</i>	5
<i>Argyrotaenia franciscana</i>	4
<i>Batia lunaris</i>	20
<i>Caloptilia stigmatella</i>	1
<i>Caloptilia syringella</i>	1
<i>Carcina quercana</i>	8
<i>Chionodes mediofuscella</i>	5
<i>Chloroclystis rectangulata</i>	1
<i>Choristoneura rosaceana</i>	11
<i>Clepsis consimilana</i>	2
<i>Clepsis virescana</i>	1
<i>Dicymolomia metalliferalis</i>	14
<i>Ditula angustiorana</i>	10
<i>Enarmonia formosana</i>	2
<i>Epinotia albangulana</i>	1
<i>Epinotia subviridis</i>	3
<i>Grapholita packardi</i>	4
<i>Grapholitha prunivora</i>	126
<i>Oegoconia quadripuncta</i>	13
<i>Recurvaria nanella</i>	17
<i>Rhopobota naevana</i>	7
<i>Setiostoma sp.</i>	1
<i>Spilonota ocellana</i>	1
<b>Total</b>	<b>293</b>

-----  
This project was a cooperative effort of the Washington State Dept. of Agriculture, Agriculture and Agri-Food Canada, and the USDA Animal and Plant Health Inspection Service (APHIS). Funding for field and lab support staff was provided in part by a Cooperative Agricultural Pest Survey (CAPS) grant from the USDA APHIS Western Region (#02-8553-0249-CA)

Eric LaGasa, Chief Entomologist  
Washington State Department of Agriculture  
Pest Program / Plant Protection Division  
P.O. Box 42560 - 1111 Washington Street  
Olympia, Washington 98504-2560  
(360) 902-2063 FAX (360) 902-2094  
elagasa@agr.wa.gov